## **Hospitalizations from falls**

Health Priority G: Intentional and Unintentional Injuries and Violence

Objective G3b: Reduce the rate of hospitalizations due to falls.

2010 Target: No target established

Indicator: Hospitalizations from falls

Wisconsin Hospitalizations due to Falls, by Sex and Ethnicity

|      |                  | Number of Hospitalizations |       |                |           |          |         |  |  |
|------|------------------|----------------------------|-------|----------------|-----------|----------|---------|--|--|
|      | Hospitalizations |                            | S     | ex             | Ethnicity |          |         |  |  |
|      | per 100,000      |                            |       |                |           | Not      |         |  |  |
| Year | Population       | Total                      | Males | <b>Females</b> | Hispanic  | Hispanic | Unknown |  |  |
| 2000 | 382.4            | 21,574                     | 7,885 | 13,689         | 168       | 19,876   | 1,530   |  |  |
| 2001 | 406.1            | 23,382                     | 9,016 | 14,366         | 214       | 21,985   | 1,183   |  |  |
| 2002 | 385.8            | 22,540                     | 8,555 | 13,985         | 211       | 21,654   | 675     |  |  |
| 2003 | 375.8            | 22,306                     | 8,512 | 13,794         | 232       | 21,462   | 612     |  |  |
| 2004 | 347.3            | 20,920                     | 7,994 | 12,926         | 238       | 20,190   | 492     |  |  |

Source: Inpatient hospitalization discharge file, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services; prepared from data collected by the Bureau of Health Information and Policy through September 30, 2003 and thereafter by the Wisconsin Hospital Association Information Center, Inc.

Note: Rates are age-adjusted to the 2000 U.S. standard population.

Wisconsin Hospitalizations due to Falls, by Race

|      | Hospitalizations | Number of Hospitalizations |           |          |        |        |        |          |  |
|------|------------------|----------------------------|-----------|----------|--------|--------|--------|----------|--|
|      | per 100,000      |                            | African   | American |        |        |        |          |  |
| Year | Population       | Total                      | American* | Indian*  | Asian* | White* | Other* | Unknown* |  |
| 2000 | 382.4            | 21,574                     | 604       | 87       | 58     | 19,481 | 232    | 1,112    |  |
| 2001 | 406.1            | 23,382                     | 637       | 106      | 65     | 21,586 | 263    | 725      |  |
| 2002 | 385.8            | 22,540                     | 598       | 134      | 52     | 21,241 | 263    | 252      |  |
| 2003 | 375.8            | 22,306                     | 641       | 112      | 61     | 21,000 | 256    | 236      |  |
| 2004 | 347.3            | 20,920                     | 583       | 109      | 62     | 19,649 | 247    | 270      |  |

Source: Inpatient hospitalization discharge file, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services; prepared from data collected by the Bureau of Health Information and Policy through September 30, 2003 and thereafter by the Wisconsin Hospital Association Information Center, Inc.

Note: Rates are age-adjusted to the 2000 U.S. standard population.

## **Continued on next page**

Prepared for Tracking the State Health Plan, 2010—State-Level Data <a href="http://dhfs.wisconsin.gov/statehealthplan/track2010">http://dhfs.wisconsin.gov/statehealthplan/track2010</a> (July 2006)
Bureau of Health Information and Policy
Division of Public Health
Wisconsin Department of Health and Family Services

<sup>\*</sup>Race groups in this table include Hispanics.

## Hospitalizations from falls, continued

Wisconsin Hospitalizations due to Falls, by Age

|      | Hospitalizations | Number of Hospitalizations |          |      |       |       |       |       |       |
|------|------------------|----------------------------|----------|------|-------|-------|-------|-------|-------|
|      | per 100,000      |                            | Age      | Ages | Ages  | Ages  | Ages  | Ages  | Ages  |
| Year | Population       | Total                      | < 1 year | 1-14 | 15-24 | 25-44 | 45-64 | 65-84 | 85+   |
| 2000 | 382.4            | 21,574                     | 79       | 692  | 513   | 1,738 | 2,960 | 9,329 | 6,263 |
| 2001 | 406.1            | 23,382                     | 77       | 751  | 596   | 2,064 | 3,576 | 9,762 | 6,556 |
| 2002 | 385.8            | 22,540                     | 71       | 732  | 556   | 1,813 | 3,312 | 9,503 | 6,553 |
| 2003 | 375.8            | 22,306                     | 58       | 679  | 526   | 1,791 | 3,436 | 9,361 | 6,455 |
| 2004 | 347.3            | 20,920                     | 70       | 681  | 594   | 1,653 | 3,463 | 8,511 | 5,948 |

Source: Inpatient hospitalization discharge file, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services; prepared from data collected by the Bureau of Health Information and Policy through September 30, 2003 and thereafter by the Wisconsin Hospital Association Information Center, Inc.

Note: Rates are age-adjusted to the 2000 U.S. standard population.